

**EC FERTILISER**  
**Zinco Solfato monoidrato 35 % Zn**  
**F.U.**  
**ZINC SALT (SULPHATE)**

**ANALYSIS**

Zinc (Zn) water soluble.....35 %

**CHARACTERISTICS AND NUTRITIONAL TARGETS**

A water soluble powder formulation based on monohydrate sulphate zinc. It prevents and cures zinc deficiency. It improves plant development as zinc is the catalytic element in the synthesis of tryptophan, amino acid precursor of indole-3-acetic acid, a natural auxin substance growth promoter.

**METHODS OF APPLICATION AND DOSAGES.**

**Zinco Solfato Monoidrato 35 % Zn F.U.** by fertigation system, mixed with other nutrients in order to create balanced solutions or on its own for zinc deficiency, with a dosage of 4-5 kg/ha. The product can be used by foliar application with a dosage of 1,5-2 kg/ha on all crops. It is suitable for soil less crops.

**PACKING:** kg 1 – kg 2,5 – kg 5 – kg 10 – kg 25

**EC FERTILISER**  
**Zinco 125 L.**  
**ZINC- BASED FERTILISER SOLUTIN (Chloride)**

**ANALYSIS**

Zinc (Zn) water soluble

p/p	p/v	g/lt
10 %	12,5 %	125

**TECHNICAL PARAMETRES**

pH of a solution at 0,1 %:	6,8-7
specific gravity at 20 °C:	1,250 kg/lt

**CHARACTERISTICS AND NUTRITIONAL TARGETS**

A liquid formulation based on zinc activated with EDTA which maintains ingredients stability even in alkaline water. Zinc is the catalytic element in the synthesis of tryptophan, amino acid precursor of indole-3-acetic acid, a natural auxin substance growth promoter.

**Zinco 125 L.** optimizes plant growth and root mass structure development.

**METHODS OF APPLICATION AND DOSAGES**

**Zinco 125 L.** can be used on all fruit trees, horticultural and ornamental crops with symptoms of underdevelopment and zinc deficiency symptoms. By foliar application, 2-3 applications with a dosage of 100-130 g/100 litres of water. As a cure, when first symptoms appear (peach trees suffering from premature leaf fall and apical leaf curling; citrus trees suffering from leaf yellowing, tomato and flower crops form browning and leaf necrosis, potato crops form brown grey and bronze coloured spots), weekly applications with a dosage of 150-180 g/100 litres of water until symptoms have completely disappeared.

The product can also be applied by fertigation system with a dosage of 10-15 kg per hectare.

**PACKING:** kg 1 – kg 5 – kg 10 – kg 25



*Special fertilisers*

**BioKimia International S.r.l.** • Via San Carlo 12/4

40023 Castel Guelfo di Bologna (BO) Italy • Tel. e Fax +39 0542 670654 • [biokimia@biokimia.it](mailto:biokimia@biokimia.it) • [www.biokimia.it](http://www.biokimia.it)